



Product Specifications

ACRYLAMIDE:BISACRYLAMIDE 19:1

<u>Catalog #</u>	<u>Size</u>
IB70022	40gm
IB70023	200gm

Physical Specifications

Acrylamide Component

CAS#	79-06-1
Formula Weight	71.08
Molecular Formula	C ₃ H ₅ NO
Purity	>99.9%
Melting Point	85 ± 1°C
pH (10%, 0.1 M NaCL)	6.5 ± 0.5
Acrylic Acid	<0.001%
Conductivity	<5µmho
Iron	<0.0001%
Lead	<0.0001%
A290 (1%,water)	<0.15

Bisacrylamide Component

CAS#	110-26-9
Formula Weight	154.17
Molecular Formula	C ₇ H ₁₀ O ₂ N ₂
Purity	>99.0%
Conductivity	<10 µmho
Acrylic Acid	<0.001%
A ₂₉₀ (2%, water)	<0.4

Molecular Biology Specifications

Dnase assay	None Detected
Rnase assay	None Detected
Protease assay	None Detected
Polymerization assay	Pass
Gel analysis	Pass

Recommended Use

A primary component of polyacrylamide gels.

To make a 30% solution:

For the 40gm size, bring the final volume to 133ml.

For the 200gm size, bring the final volume to 667ml.

To make a 40% solution:

For the 40gm size, bring the final volume to 100ml.

For the 200gm size, bring the final volume to 500ml.

Storage

Store at room temperature. Keep tightly sealed. Protect from moisture.

Warning

Poison/Neurotoxin/Lachrymator. Suspected carcinogen. May be harmful or fatal if swallowed. Avoid breathing dust. Avoid contact with eyes, skin, and clothing. Wash thoroughly with water after handling. See Material Safety Data Sheet for additional information.

FOR RESEARCH AND DEVELOPMENT PURPOSES ONLY

5/7/2009